Rehabilitation Protocol: Medial Patellofemoral Ligament Reconstruction

OMAR RAHMAN, MD, MBA

Orthopedic Surgeon | Sports Medicine

ARTHROSCOPIC SURGERY | JOINT RECONSTRUCTION



Name:	Date:
Diagnosis:	Date of Surgery:
Phase I (Weeks 0-2) Weightbearing: Toe-touch weightbearing with cre Hinged Knee Brace Cocked in full extension for ambulation and Range of Motion – PROM/AAROM Range of Motion – PROM/AAROM Therapeutic Exercises Quad/Hamstring sets Heel slides/Prone hangs/Patellar mobilization of Straight-Leg Raise with brace in full extensions.	d sleeping
Phase II (Weeks 2-6) ■ Weightbearing: As tolerated – wean from crutch is a Hinged Knee Brace: Unlocked ■ Range of Motion – PROM/AAROM/AROM: Maintain flexion (goal of 90 degrees by week 6) ■ Therapeutic Exercises ■ LIMIT WEIGHBEARING EXERCISE TO FLEX is Isometric quadriceps strengthening ■ Heel slides/Prone hangs ■ Patellar mobilization	ain full knee extension – work on progressive knee
Phase III (Weeks 6-12) Weightbearing: Full weightbearing without crutce Discontinue Knee Brace - Convert to Pa Range of Motion - Advance to Full/Painless ROM Therapeutic Exercises Wall sits/Lunges Proprioception training Stationary bicycle	itellar Stabilization Sleeve (Reaction Brace)
 Phase IV (Months 3-4) Advance closed chain strengthening – leg press, leg Plyometric and proprioception training Treadmill jogging/Elliptical 	g curls
 Phase V (Months 4-6) Gradual return to athletic activity as tolerated Maintenance program for strength and endurance 	•
Comments:	
Frequency: times per week	Duration: weeks