Pectoralis Major Repair Post-Operative Rehabilitation Protocol

OMAR RAHMAN, MD, MBA

Orthopedic Surgeon | Sports Medicine





Name:	Date:
Diagnosis	SEPISEPISEP SEPISEPISEPISEPIDate SEP Of SEP Surgery: SEP
PTPT SEP!SEP!	SEPISEPISEPISEPIDATESEPSOTISEPSOURGERY: SEPI
SIRaTI	ng Immobilizer: Worn at all times – Sleep with pillow under elbow to support the operative arm nge of Motion: Supported pendulum exercises under guidance of PT erapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side), and exercises, patient may ride stationary bike with operative arm in the sling
SIRaT	It isp 4 - 6 p Weeks sp and Immobilizer: Worn at all times – Sleep with pillow under elbow to support the operative arm nge of Motion: AAROM in the supine position with wand – Goal: Forward Flexion to 90 degrees erapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side), and exercises, shoulder shrugs/scapular retraction without resistance
SIRa	III: [F] 6 - 8 SEE Weeks Immobilizer: May be discontinued Immobilizer: M
RateT	record of Motion: AROM and AAROM in the pain free range – No PROM, Goals: Full ROM erapeutic Exercises: Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone apular Retraction Exercises (without weights), Wall Push-Ups (no elbow flexion > 90 degrees)
RaTl	N: [P] 3 - 6 [F] Months nge of Motion: Full ROM erapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May star th weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups turn to athletics at 6 months post-op
Comment	[P] [SEP]
Frequenc	SEPSEP SEP TIMES SEP Duration: SEPSEP SEP SEP SEP SEP SEP SEP SEP SEP
Sionaturo	ELE Date: